



Rodenticide Comparison Chart



Active Ingredient	Brand					Type	Notes
	Liphatech	Neogen	Motomco	Scimetrics	Others		
Brodifacoum		Havoc	JAGUAR		Weatherblok XT	Anticoagulant	2nd generation. Death occurs approximately 4-5 days after lethal consumption
Difenacoum		DI-Kill				Anticoagulant	2nd generation. Death occurs approximately 4-5 days after lethal consumption
Bromadiolone	Boothill Revolver SOFT BAIT		HAWK		Contract ALL-WEATHER BRIGAND RODENTICIDES	Anticoagulant	2nd generation. 4-5 days post feeding
Difethialone	HOMBRE FASTDRAW FastDraw takes 4-5 days to work.					Anticoagulant	2nd generation. 4-5 days post feeding
Chlorophacinone	rozol					Anticoagulant	1st generation. 5-7 days post feeding
Diphacinone		Ramik	Toucat		JT EATON BAIT BLOCK	Anticoagulant	1st generation. Death occurs approximately 4-5 days after lethal consumption
Warfarin		Rodex		Kaput		Anticoagulant	1st generation. Death occurs approximately 4-5 days after lethal consumption
Bromethaline	GUNSLINGER	CyKill	RAMPAGE			Neurotoxicant	Acute toxicant. 2+ days post feeding
Cholecalciferol			Agrid₃ BAIT CHUNX			Vitamin D	Acute toxicant. 5 days post feeding; For organic production.
Zinc Phosphide		PROZAP			ZP TRACKING POWDER	Systemic Toxicant	Acute toxicant. Death may occur 2 days after lethal consumption.

Iowa	Minnesota	Indiana	Delaware
1716 Detroit St Ellsworth, IA 50075 888-378-4045	325 Lakeland Dr NE Willmar, MN 56201 800-533-1899	1854 East Waller Dr Washington, IN 47501 877-254-3410	31024 Country Gardens Blvd #G2 Dagsboro, DE 19939 877-732-3894

Restricted Use